



US006372279B1

(12) **United States Patent**
Ishida et al.

(10) Patent No.: **US 6,372,279 B1**
(45) Date of Patent: **Apr. 16, 2002**

(54) **SWEETENER COMPOSITION**

(75) Inventors: **Hirotoishi Ishida; Akihiro Kishishita;**
Takeshi Nagai; Kazutaka Nagashima;
Atsuhiko Hirano, all of Kawasaki (JP)

(73) Assignee: **Ajinomoto Co., Inc., Toyko (JP)**

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **09/707,954**

(22) Filed: **Nov. 8, 2000**

Related U.S. Application Data

(63) Continuation of application No. PCT/JP99/02197, filed on Apr. 26, 1999.

(30) Foreign Application Priority Data

May 8, 1998 (JP) 10-125990

(51) Int. Cl.⁷ **A23L 1/236**

(52) U.S. Cl. **426/548; 426/590; 560/40**

(58) Field of Search **426/548, 590;**
560/39, 40, 41

(56) References Cited

U.S. PATENT DOCUMENTS

5,480,668 A * 1/1996 Nofre et al. 426/548
6,048,999 A * 4/2000 Pajor et al. 560/39

FOREIGN PATENT DOCUMENTS

JP 8-503206 4/1996
JP 10-248521 9/1998

* cited by examiner

Primary Examiner—Leslie Wong

(74) Attorney, Agent, or Firm—Oblon, Spivak, McClelland, Maier & Neustadt, P.C.

(57) ABSTRACT

One embodiment of the present invention provides a sweetener composition, which includes a mixture of N-{N-(3,3-dimethylbutyl)-L- α -aspartyl}-L-phenylalanine 1-methyl ester, and Acesulfame K, wherein a ratio of the Acesulfame K to a total amount of the N-{N-(3,3-dimethylbutyl)-L- α -aspartyl}-L-phenylalanine 1-methyl ester and the Acesulfame K is in the range of 10 to 97% by weight, methods of making and of using. Another embodiment of the present invention provides a method for preparing a sweetener composition, which includes drying an A-type crystal of N-{N-(3,3-dimethylbutyl)-L- α -aspartyl}-L-phenylalanine 1-methyl ester to obtain a C-type crystal of N-{N-(3,3-dimethylbutyl)-L- α -aspartyl}-L-phenylalanine 1-methyl ester. Another embodiment of the present invention provides a method for producing a sweetener, which includes admixing N-{N-(3,3-dimethylbutyl)-L- α -aspartyl}-L-phenylalanine 1-methyl ester with Acesulfame K, wherein a ratio of the Acesulfame K to a total amount of the N-{N-(3,3-dimethylbutyl)-L- α -aspartyl}-L-phenylalanine 1-methyl ester and the Acesulfame K is in the range of 10 to 97% by weight. Another embodiment of the present invention provides a method for improving the dissolution rate of N-{N-(3,3-dimethylbutyl)-L- α -aspartyl}-L-phenylalanine 1-methyl ester, which includes, prior to dissolving the N-{N-(3,3-dimethylbutyl)-L- α -aspartyl}-L-phenylalanine 1-methyl ester, admixing the N-{N-(3,3-dimethylbutyl)-L- α -aspartyl}-L-phenylalanine 1-methyl ester with Acesulfame K, wherein a ratio of the Acesulfame K to a total amount of the N-{N-(3,3-dimethylbutyl)-L- α -aspartyl}-L-phenylalanine 1-methyl ester and the Acesulfame K is in the range of 10 to 97% by weight.

17 Claims, 2 Drawing Sheets